

# User's Guide for CASA2WRF

Zhining Tao, Jossy P. Jacob and Eric Kemp

July 21, 2014

## Contents

<b>1</b>	<b>Overview</b>	<b>2</b>
<b>2</b>	<b>Using the Software</b>	<b>2</b>
2.1	CASACO2 pre-processor . . . . .	3
2.2	Run CASA2WRF . . . . .	3
2.3	Run WRFCHM . . . . .	4
<b>3</b>	<b>References</b>	<b>4</b>

## 1 Overview

Coupled NU-WRF-CASA modeling system has the capability for tracer CO<sub>2</sub> (i.e., no interaction with chemistry) simulations. It is expected to resolve small scale CO<sub>2</sub> sources and sinks, reduce transport uncertainties, and contribute to improving global CO<sub>2</sub> modeling. This requires specifying initial, lateral boundary, and flux emission fields of CO<sub>2</sub>. To that end, several utilities have been developed to process the Carnegie-Ames-Stanford-Approach (CASA) global CO<sub>2</sub> concentrations and fluxes and provide them to NU-WRF.

CASA CO<sub>2</sub> concentrations are based on the Goddard Space Flight Center Parameterized Chemistry and Transport Model (PCTM), which is driven by the real-time meteorological fields from the Goddard Global Modeling and Assimilation Office, version 5 (GEOS-5). The biospheric CO<sub>2</sub> fluxes are produced from CASA, the biomass burning emissions are from the Global Fire Emissions Database (GFED), and the oceanic and anthropogenic CO<sub>2</sub> fluxes are based on the sources described by Kawa et al.[2004, 2010]. The hourly model output is at the resolution of 1 x 1.25 degree with 28 vertical levels. The PCTM/GEOS5/CASA-GFED model has been widely tested, and has shown favorable results in carbon cycle comparison studies [e.g., Kawa et al., 2010, and references therein].

CASA2WRF is the NU-WRF utility for preprocessing the CO<sub>2</sub> concentration and flux data to furnish initial and boundary conditions to WRF. It reads the CASA CO<sub>2</sub> concentration in the NETCDF format, interpolates it to WRF domains (single or nested), and appends variable `?casaco2?` to `wrfinput` and `wrfbdy` files at given time intervals.. Additionally, CASA2WRF utility reads the CO<sub>2</sub> flux data, interpolate them to WRF domains (single or nested), and write to NETCDF files readable to NU-WRF with a frequency specified in the WRF namelist file. The capability to interpolate the flux data at each simulation time step in NU-WRF is achieved by adding the flux tendency (rate of flux change) to the flux data files.

## 2 Using the Software

To compile CASA2WRF, the user must type `./build.sh casa2wrf` or `./build.sh allchem` and executables for pre-processing CASA CO<sub>2</sub> data and `casa2wrf` will be created. The workflow for using CASACO2 is listed below.

1. Run GEOGRID
2. Run UNGRIB (or MERRA2WRF or GEOS2WRF)
3. Run METGRID
4. Run REAL
5. Run CASA2WRF

## 6. Run WRF-Chem with CASA option

There are 3 steps for including CASACO2 in NU-WRF.

### 2.1 CASACO2 pre-processor

Pre-processor for CASA2WRF (`$NUWRFDIR/Utils/casa2wrf/pproc/`) converts the binary input files from PCTM to NETCDF format and add a timestamp in the WRF time format.

- CO<sub>2</sub> Concentration data: Compile with `./build.sh casa2wrf` or to compile separately: `make Makefile_CO2_conc` (or compile with NUWRF `build.sh casa2wrf`) and to run: `./Read_CO2_conc.x filename indir`; creates netcdf files in `conc/CASACO2.*.nc`
- CO<sub>2</sub> flux data: Compile with NUWRF `build.sh casa2wrf` or to compile separately: `make -f Makefile_CO2_Flux` and to run: `Read_CO2_Flux.x filename indir`; creates yearly NETCDF data file: `flux/CO2flux *.nc`

### 2.2 Run CASA2WRF

- Compile: `./build.sh casa2wrf`
- Run: `./casa2wrf` or use batch scripts: `run_casa2wrf_discover.sh`.
- Make sure that the `wrfinput_d*` and `wrfbdy_d01` exists in your `rundir`.
- Output: `wrfinput` and `wrfbdy` files will be modified, and flux datafiles will be generated in `chem_flux/` directory.

The `namelist.casa2wrf` contains the following information:

Variable Names	Description
<code>&amp;wrf</code>	
<code>max_dom</code>	integer, specifies number of domains.
<code>wrf_dir</code>	string, WRF run directory
<code>flux_only</code>	integer, =0 for processing CO2 concentration and flux data, =1 for processing CO2 flux emission data only.
<code>fluxdt</code>	real, time interval of input flux emission data in hours.
<code>&amp;casa_conc</code>	
<code>casa_format</code>	integer, =5 for netcdf files
<code>casa_dir</code>	string, CASACO2 concentration data directory
<code>casa_prefix</code>	string, CASACO2 concentration datafile prefix
<code>&amp;casa_flux</code>	
<code>casa_format</code>	integer, =5 for netcdf files
<code>casa_dir</code>	string, CASACO2 flux emission data directory
<code>casa_prefix</code>	string, CASACO2 flux datafile prefix

## 2.3 Run WRF-CHEM

To run WRF-Chem with CASACO2, the **namelist.input** file should have the following information:

Variable Names	Description
&time_control	
auxinput18_inname	string, = "chem_flux/CO2_(domain)_(date)", flux datafile name
auxinput18_interval_m	integer, time interval of input flux data in minutes for each domain e.g. for 3 hourly input data = 180,180,
FRAMES_PER_auxinput18	integer, number of dataset in each datafile e.g. for 1 data set in each file/ 2 domains = 1,1
IO_FORM_HISTORY	integer, = 2, for netcdf file
IO_FORM_RESTART	integer, = 2, for netcdf file
IO_FORM_INPUT	integer, = 2, for netcdf input file
IO_FORM_BOUNDARY	integer, = 2, for netcdf file
io_form_auxinput18	integer, = 2, for netcdf input file
&chem	
chem_opt	integer, for casaco2, chem_opt for each domain =18, 18,
io_style_emissions	integer, emission input file format = 2 for netcdf
casafxdt	emission data interval in minutes for each domain, e.g. for 3 hourly files casafxdt = 180., 180.,
emiss_inpt_opt	integer, for casaco2 case = 18, 18,
emiss_opt	integer, for casaco2 case = 18, 18,
chem_in_opt	integer, = 1, 1,
emi_inname	string, emission input filename, e.g. = "chem_flux/CO2_"
emiss_opt_vol	integer, = 0,0,
phot_opt	integer, = 2, 2,
gas_drydep_opt	integer, = 0, 0,
gas_bc_opt	integer, for casaco2 case = 18, 18,
gas_ic_opt	integer, for casaco2 case = 18, 18,
have_bcs_chem	logical, Lateral boundary condition is provided for outer domain only e.g. = .true., .false.,

## 3 References

Kawa, S. R., D. J. Erickson III, S. Pawson, and Z. Zhu (2004), Global CO2 transport simulations using meteorological data from the NASA data assimilation system, *J. Geophys. Res.*,109, D18312, doi:10.1029/2004JD004554.

Kawa, S. R., J. Mao, J. B. Abshire, G. J. Collatz, X. Sun, and C. J. Weaver (2010), Simulation studies for a space-based CO2 lidar mission, *Tellus, Ser. B*, 62 (5), 759 ? 769, doi:10.1111/j.1600-0889.2010.00486.x